

学位論文

**Current status and management of pancreatic trauma with
main pancreatic duct injury: A multicenter nationwide survey
in Japan**

香川大学大学院医学系研究科

医学 専攻

安藤 恭久



ORIGINAL ARTICLE

Current status and management of pancreatic trauma with main pancreatic duct injury: A multicenter nationwide survey in Japan

Yasuhisa Ando¹ | Keiichi Okano¹ | Hiroshi Yasumatsu³ | Toshimasa Okada⁴ | Kimiyoshi Mizunuma⁵ | Minoru Takada⁶ | Shinjiro Kobayashi⁷ | Keisuke Suzuki⁸ | Nobuya Kitamura⁹ | Minoru Oshima¹ | Hironobu Suto¹ | Miyatake Nobuyuki² | Yasuyuki Suzuki¹

¹Department of Gastroenterological Surgery, Kagawa University, Kita-gun, Japan

²Department of Hygiene, Kagawa University, Kita-gun, Japan

³Shock and Trauma Center/Hokusoh HEMS, Nippon Medical School Chiba Hokusoh Hospital, Inzai, Japan

⁴Department of Digestive Surgery, Kawasaki Medical School, Kurashiki-City, Japan

⁵Department of Radiology, Nasu Red Cross Hospital, Otawara-City, Japan

⁶Department of Surgery, Teine Keijinkai Hospital, Sapporo-City, Japan

⁷Division of Gastroenterological and General Surgery, St. Marianna University School of Medicine, Kawasaki-City, Japan

⁸Trauma and Acute Critical Care Center, Tokyo Medical and Dental University Hospital of Medicine, Bunkyo-ku, Japan

⁹Department of Emergency and Critical Care Medicine, Kimitsu Chuo Hospital, Kisarazu-City, Japan

Correspondence

Yasuhisa Ando, Department of Gastroenterological Surgery, Kagawa University, 1750-1 Ikenobe, Miki-cho, Kita-gun, Kagawa 761-0793, Japan.
Email: y-ando@med.kagawa-u.ac.jp

Funding information

Project study funds of Japanese Society for Abdominal Emergency Medicine, Grant/Award Number: 2016-P-02

Abstract

Background: Pancreatic trauma is reportedly associated with high morbidity and mortality. Main pancreatic duct (MPD) injury is critical for treatment.

Methods: As a study project of the Japanese Society for Abdominal Emergency Medicine (JSAEM), we collected the data of 163 patients with pancreatic trauma who were diagnosed and treated at JSAEM board-certified hospitals from 2006 to 2016. Clinical backgrounds, diagnostic approaches, management strategies, and outcomes were evaluated.

Results: Sixty-four patients (39%) were diagnosed as having pancreatic trauma with MPD injury that resulted in 3% mortality. Blunt trauma and isolated pancreatic injury were independent factors predicting MPD injury. Nine of 11 patients with MPD injury who were initially treated nonoperatively had serious clinical sequelae and five (45%) required surgery as a secondary treatment. Among all cases, the detectability of MPD injury of endoscopic retrograde pancreatography (ERP) was superior to that of other imaging modalities (CT or MRI), with higher sensitivity and specificity (sensitivity = 1.0; specificity = 0.93).

Conclusions: Acceptable outcomes were observed in pancreatic trauma patients with MPD injury. Nonoperative management should be carefully selected for MPD injury. ERP is recommended to be performed in patients with suspected MPD injury and stable hemodynamics.

KEY WORDS

endoscopic retrograde pancreatography, main pancreatic duct injury, management, nationwide survey, pancreatic trauma

Yasuhisa Ando and Keiichi Okano contributed equally to this work.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2020 The Authors. *Journal of Hepato-Biliary-Pancreatic Sciences* published by John Wiley & Sons Australia, Ltd on behalf of Japanese Society of Hepato-Biliary-Pancreatic Surgery.

1 | INTRODUCTION

Pancreatic trauma is relatively rare, with an overall frequency of only 0.2%-0.3% among all trauma but a high mortality rate of 17%-46%.¹⁻⁴ Pancreatic fistula is one of the major complications of pancreatic trauma. This can contribute to prolonged hospital stays and fatal outcomes. Some reports stated that delayed diagnosis and inappropriate treatment of main pancreatic duct (MPD) injury could increase both complications and mortality rates.⁴⁻⁶ However, because of limited cases at a single institution, experienced specialists and high-quality evidence are scarce. Consequently, a consensus regarding diagnostic and treatment algorithms as well as optimal surgical selection has not been established.

Operative management (OM) is often selected for deep injury, especially severe pancreatic trauma with MPD injury. Pancreatic-sparing surgery may be selected depending on the patient's age and general condition, the extent of injury, and the skill of the surgeon. In recent years, some reports have recommended conservative pancreatic drainage for pancreatic trauma with MPD injury, but the short- and long-term outcomes remain unclear.⁶⁻⁸ Because accurate evaluation of MPD injury by abdominal computed tomography (CT) scan alone is often difficult, some reports recommended emergency endoscopic retrograde pancreatography (ERP) if the patient's general condition is stable.⁶⁻⁸ In some cases with MPD injury, nonoperative management (NOM) with endoscopic pancreatic stenting (EPS) is possible after diagnosis by ERP; however, comprehensive evaluation of ERP for pancreatic trauma has not yet been performed.

This study aimed to evaluate the current status and optimal management strategy, including the diagnostic role of ERP, for pancreatic trauma by conducting a multicenter nationwide survey in Japan. In particular, we focused more on cases with MPD injuries that are difficult to diagnose and treat.

2 | METHODS

2.1 | Patients and data collection

We conducted a questionnaire survey at abdominal emergency specialist institutions as a project study of the Japanese Society for Abdominal Emergency Medicine (JSAEM). Clinical data were collected from 173 patients who were diagnosed and treated for pancreatic trauma at the 31 board-certified hospitals in JSAEM between January 2006 and December 2016. Furthermore, 10 patients with cardiopulmonary arrest before or immediately after arrival who died within 24 hours after examination or were transferred to another hospital for treatment were excluded. We retrospectively analyzed patients' demographics, physical findings,

and laboratory data within 48 hours after injury ($n = 144$), along with existence of the MPD injury, imaging modalities, treatment strategies (NOM/OM), perioperative variables, and outcomes in the finally enrolled 163 patients.

2.2 | Groups and definitions

All pancreatic trauma cases were divided into the groups of MPD intact and MPD injured according to the final diagnosis made in each center, being judged by one or some imaging modalities combined with clinical, intraoperative and/or pathological findings. Major complications associated with treatment were defined as those with a Clavien–Dindo classification⁹ of \geq IIIa. A postoperative pancreatic fistula (POPF) was defined according to the criteria of the International Study Group on Pancreatic Fistula (ISGPF).¹⁰

2.3 | Statistical analysis

Descriptive statistics are reported as mean (standard deviation) for normally distributed continuous variables, median (range) for other continuous variables, and frequency (percent) for categorical variables. Chi-squared test or Fischer exact test was used to analyze categorical variables of two groups, whereas unpaired *t*-test or Mann–Whitney *U*-test was used to compare continuous variables. Additionally, analysis of covariance was used to adjust for confounding factors. The level of significance was set at $P < .05$. JMP Pro version 14 (SAS Institute Inc.) was used to conduct all statistical analyses.

2.4 | Ethical statements

The protocol for this research project has been approved by the Committee of JSAEM (Approval No. 18-1-R1), and the Committee of Kagawa University (Approval No. 29-200). It conforms to the provisions of the Declaration of Helsinki. Consent was obtained from the subjects in an informative or disclosure document.

3 | RESULTS

3.1 | Patient characteristics

A total of 163 cases were classified into two groups, "MPD intact" ($n = 99$) and "MPD injured" ($n = 64$). Table 1 shows the clinical characteristics of patients divided into these groups. There were 109 (67%) males and 54 (33%) females, with a mean age of 42.7 years. As indicated in Figure S1,

TABLE 1 Demographics and clinical characteristics of patients with or without main pancreatic duct (MPD) injury (number [%] or value)

Variables	Overall n = 163	MPD intact n = 99	MPD injured n = 64	Univariate analysis	Multivariate analysis
Gender					
Male/Female	109(67)/54 (33)	62 (63)/37 (37)	47 (73)/17 (27)	0.152	
Age (y)	42.7 ± 1.8	45.6 ± 2.3	38.3 ± 2.8	0.045	0.388
Cause of trauma					
Traffic accident	90 (55)	55 (56)	35 (55)	0.913	
Non-traffic accident	73 (45)	44 (44)	29 (45)		
Injury type					
Blunt	140 (86)	78 (79)	62 (97)	0.001	0.007
Penetrating	23 (14)	21 (21)	2 (3)		
Intraabdominal injury					
Pancreas only	55 (34)	23 (23)	32 (50)	<0.001	0.013
Pancreas injury location					
Head	66 (40)	47 (47)	19 (30)	0.033	
Body or tail	97 (60)	52 (53)	45 (70)		
Injury Severity Score (ISS)	15.8 ± 0.9	16.2 ± 1.1	15.3 ± 1.4	0.613	0.998
Revised Trauma Score (RTS)	7.3 ± 0.1	7.3 ± 0.1	7.4 ± 0.1	0.655	

Note: Frequency (%) for categorical variables.

Mean (standard deviation [SD]) for continuous variables.

male patients showed a bimodal distribution and tended to be in their 10s and 60s. In contrast, female patients tended to be young, especially in their teens. As presented in Table 1, there were 140 patients (86%) with blunt trauma and 90 (55%) with injury caused by traffic accidents. Fifty-five patients (34%) had isolated pancreatic injury, and 108 (66%) were associated with coexisting severe abdominal injury. Blunt trauma and isolated pancreatic injury were revealed as the independent clinical factors predicting MPD injury when adjusted by multivariate analysis ($P = .007$ and $P = .013$, respectively). Concurrent liver damage was most frequently

observed in 24 patients (15%). Major vascular damage, considered fatal, occurred in 13 patients (8%). Anatomical injury sites included the head of the pancreas in 66 patients (40%) and the pancreatic body/tail in 97 (60%). The mean injury severity score (ISS) was 15.8. The mean serum amylase level determined within 48 hours after injury was 332.4 U/L. Figure 1 shows laboratory values within 48 hours of injury between the groups with and without MPD injury. Serum amylase levels as well as CRP were significantly higher in patients with MPD injury compared to those without MPD injury (506.1 ± 64 vs 220.0 ± 51 , $P < .001$).




FIGURE 1 Laboratory values within 48 h of injury between the groups with and without MPD injury (n = 144). Serum amylase levels in the vertical axis are displayed using a logarithmic scale. WBC (μL ; $13\,491 \pm 609$ vs $13\,328 \pm 749$, $P = .865$). CRP (mg/dL ; 0.56 ± 0.5 vs 2.44 ± 51 , $P = .010$). Amylase (U/L ; 220.0 ± 51 vs 506.1 ± 64 , $P < .001$). CRP, C-reactive protein; MPD, main pancreatic duct; WBC, white blood cell. One-way analysis of variance

3.2 | Treatment and outcomes

Treatment and outcomes of the patients are shown in Table 2. Forty-one (25%) patients underwent ERP; significantly more patients with MPD injury underwent ERP than those with MPD intact (14 [14%] vs 27[42%]). Furthermore, 58 (36%) patients were nonoperatively treated, whereas 105 (64%) were surgically treated. In addition, 53 (83%) patients with MPD injury and 52 (53%) with MPD intact had OM ($P < .001$). Clinical sequelae were observed significantly more frequently in patients with MPD injury, particularly in the NOM group. Moreover, 17 (29%) and 49 (47%) patients had one or more sequential problems in NOM and OM groups, respectively. The average length of stay in hospital was 43.2 days (37.6 and 51.8 days in MPD intact and MPD injured cases, respectively, $P = .055$). There were eight (5%) hospital deaths, and all but one patient died of multiple organ injury.

3.3 | Diagnostic imaging of MPD injury

Table 3 shows the imaging modalities performed to detect MPD injury in all patients. The rates of performance of CT, ERP, and magnetic resonance perfusion imaging (MRP) were 94%, 25%, and 12%, respectively. CT was performed in the majority of patients with pancreatic injury, whereas MRP was performed at a much lower rate. The sensitivities/specificities of MPD injury detection by CT, ERP, and MRP were 0.98/0.88, 1.0/0.93, and 0.89/0.8, respectively. The diagnostic accuracy of ERP was superior to that of other imaging modalities (CT and MRP) for the diagnosis of MPD injury.

3.4 | Management of MPD injury

Table 4 focuses on 64 patients with MPD injury and compares backgrounds and outcomes between their initial treatment

TABLE 2 Treatment and outcomes of patients with or without main pancreatic duct (MPD) injury (number (%) or value)

Variables	Overall n = 163	MPD intact n = 99	MPD injured n = 64	P-value
Treatment				
Rate of ERP	41 (25)	14 (14)	27 (42)	<.001
Initial treatment strategy				
Non-operative management (NOM)	58 (36)	47 (47)	11 (17)	<.001
Operative management (OM)	105 (64)	52 (53)	53 (83)	
Outcomes				
Complications ^a	66 (40)	28 (28)	38 (59)	<.001
NOM	17 (29)	8 (17)	9 (82)	<.001
OM	49 (47)	20 (38)	29 (55)	.095
Hospital stay (d)	43.2 ± 3.6	37.6 ± 4.6	51.8 ± 5.7	.055
Mortality	8 (5)	6 (6)	2 (3)	.397

Note: Frequency (%) for categorical variables.

Mean (standard deviation [SD]) for continuous variables.

Abbreviation: ERP, endoscopic retrograde pancreatography.

^aClavien–Dindo classification of grade IIIa or higher.

TABLE 3 Accuracy of imaging modalities to detect MPD injury and diagnose the type of injury in grade III patients

Imaging modality	n (%)	CT		ERP		MRP	
Number of patients	n = 84	80 (95)		35 (42)		11 (13)	
MPD injury							
Grade		Positive	Negative	Positive	Negative	Positive	Negative
IIIa		1	17	0	8	0	1
IIIb		50	12	26	1	8	2
Sensitivity		0.81		0.96		0.8	
Specificity		0.94		1.0		1.0	

Abbreviations: MPD, Main pancreatic duct; CT, Computed tomography; ERP, Endoscopic retrograde pancreatography; MRP, Magnetic resonance pancreatography.

TABLE 4 Demographics and clinical characteristics of main pancreatic duct injured patients initially treated with operative and nonoperative management

Variables	Nonoperative management n = 11	Operative management n = 53	P value
Background			
Age (y)	49.5 ± 6.9	35.9 ± 3.1	.079
Injury type			
Blunt	11 (100)	51 (96)	.513
Penetrating	0 (0)	2 (4)	
Intraabdominal injury			
Pancreas only	5 (45)	27 (51)	.740
Pancreas injury location			
Head	4 (36)	15 (28)	.594
Body or tail	7 (64)	38 (72)	
Rate of comorbidities	5 (45)	10 (23)	.058
Injury Severity Score (ISS)	14.7 ± 2.5	15.4 ± 1.1	.814
Laboratory values			
WBC (/μL)	12,825 ± 1634	13,577 ± 744	.677
CRP (mg/dL)	2.9 ± 2.2	3.2 ± 1.0	.895
Amylase (U/L)	562.8 ± 205.7	482.6 ± 91.1	.723
Treatment			
Rate of ERP	6 (55)	22 (42)	.428
Outcomes			
Complications ^a	9 (82)	29 (55)	.096
Pseudocyst ^b	4 (36)	6 (11)	
Pancreatic fistula ^b	4 (36)	22 (42)	
Abscess ^b	2 (18)	10 (19)	
Peritonitis ^b	2 (18)	2 (4)	
Conversion to surgery	5 (45)		
Hospital stay (d)	77.3 ± 13.0	47.2 ± 5.7	.038
Mortality	0 (0)	2 (4)	.513

Note: Frequency (%) for categorical variables.

Mean (standard deviation [SD]) for continuous variables.

Abbreviations: CRP, C-reactive protein; ERP, endoscopic retrograde pancreatography; WBC, white blood cell.

^aClavien–Dindo classification of grade IIIa or higher.

^bOverlapping distribution.

strategies. There were 11 patients in the NOM group and 53 in the OM group. There were no significant differences in clinical factors such as age, morphology of pancreatic injury, ISS, or blood test results at the time of visit. However, the length of hospital stay was significantly extended in the NOM group (77.3 and 47.2 days in NOM and OM groups, respectively, $P = .038$).

Table 5 lists individual data of 11 patients with MPD injury who were initially treated nonoperatively. The initial treatment included drainage in five patients, EPS in five, and octreotide injection in one. Only two (18%) patients successfully recovered after the initial treatment. In contrast, nine (82%) patients had clinical sequelae.

Furthermore, five (45%) of these nine patients required surgery as a secondary treatment. Four patients (36%) required long-term hospitalization for 90 days or more, although there was no hospital death.

Conversely, in a series of 53 patients with MPD injury who were treated with OM, 48 (90%) received pancreatic resection and five (10%) received drainage alone and/or pancreatic suture repair without pancreatic resection (Table 6). When comparing pancreatic resection and no pancreatic resection, the length of hospital stay was significantly longer in patients without pancreatic resection ($P = .003$). Patients who underwent the Letton–Wilson procedure were significantly younger ($P = .038$) and had more blood loss

TABLE 5 Individual characteristics and outcomes of main pancreatic duct (MPD) injured patients initially treated with nonoperative management

Case	Age/ Gender	Pancreas injury (location)	Type of MPD injury (CT/ERP/ MRP)	ERP	Initial treatment			Second treatment			Hospital stay (days)
					Treatment modality	Post- traumatic hours	Outcome and complications	Conversion modality	Post- traumatic days	Outcome and complications	
1	77/F	Body/Tail	Partial (CT)	No	Drainage	2	Recovered	None	None	None	235
2	61/M	Body	Complete (CT/ERP)	Yes	Drainage	1	Pseudocyst	Operation (DP)	18	Pancreatic fistula	119
3	86/M	Body	Partial (CT/ MRP)	No	Drainage	792	Pseudocyst	ENPD	unknown	Recovered	20
4	73/M	Tail	Partial (CT)	No	Drainage	1	Abscess Pancreatic fistula	ENPD	>30	Recovered	185
5	60/M	Body	Partial (CT)	No	Drainage	144	Pancreatic fistula	Operation (DP)	32	Pancreatic fistula	87
6	24/M	Head	Complete (CT/ERP)	Yes	EPS	10	Recovered	None	None	None	34
7	8/M	Head / Body	Complete (CT/ERP)	Yes	EPS	5	Peritonitis	Operation (PD)	2	Bile leakage	26
8	71/M	Head	Partial (CT/ ERP)	Yes	EPS	4	Peritonitis	Operation (Operative drainage)	2.5	Intestinal perforation	34
9	57/M	Tail	Partial (CT/ ERP)	Yes	EPS	1.5	Pseudocyst Pancreatic fistula Abscess	Drainage	4	Recovered	105
10	11/M	Head	Complete (CT/ERP/ MRP)	Yes	EPS	30	Pseudocyst	None	None	None	82
11	16/M	Body	Complete (CT/ERP/ MRP)	No	Octreotide	1	Pancreatic fistula	Operation (L&W)	24	Recovered	31

Abbreviations: CT, computed tomography; DP, distal pancreatectomy; Drainage, percutaneous drainage; ENPD, endoscopic nasopancreatic drainage; EPS, endoscopic pancreatic stenting; ERP, endoscopic retrograde pancreatography; L&W, Letton–Wilson's procedure; MRP, magnetic resonance pancreatography; PD, pancreatoduodenectomy.

($P = .010$) than those who underwent pancreaticoduodenectomy (PD). Moreover, regarding deep pancreatic body/tail injury, the Warshaw procedure was performed at a younger age ($P = .018$) than distal pancreatectomy. There were no statistically significant differences in postoperative morbidity, grade B and C POPF, and mortality rates among surgical procedures. Two patients (4%) had death outcomes secondary to hemorrhagic shock.

4 | DISCUSSION

Traumatic pancreatic injury is a relatively rare trauma but is associated with a high mortality rate.¹⁻⁴ The Japanese

trauma data bank previously reported that this injury accounted for 0.25% of all trauma, 2.4% of abdominal trauma, and 4.3% of mortality.⁴ In Japan, more than 90% of these injuries are caused by blunt trauma resulting from traffic accidents.^{4,11} MPD involvement is the most important factor of patient outcomes, and delays in diagnosis are associated with high postoperative morbidity and mortality rates, particularly in cases of MPD injury.¹² Because the pancreas is surrounded by the liver, biliary tract, spleen, stomach, duodenum, colon, and large blood vessels, the frequency of concomitant other organ injury is also high. The mortality rate is relatively high in such cases.^{4,13} The frequency of single pancreatic injury is less than 10%. The number of concurrently damaged organs averages 3.5-4.1, with

TABLE 6 Clinical characteristics and outcomes of 53 main pancreatic duct injured patients initially treated with operative management

Operation	Patient number (%)	Age (year)	Operation time (min)	Amount of bleeding (mL)	POPF \geq grade B n (%)	Postoperative complication ^a	Hospital stay (days)	Mortality n (%)
Pancreatic resection	48 (90)	29 [4-83]	258 [85-690]	1,090 [0-5831]	19 (40)	30 (63)	37 [2-130]	2 (4)
Pancreaticoduodenectomy	10 (19)	29.5 [16-77]	347.5 [85-550]	2,525.5 [563-4300]	4 (40)	5 (50)	43.5 [25-126]	0 (0)
Distal pancreatectomy	14 (26)	42 [11-78]	176.5 [85-360]	1908 [100-5831]	6 (43)	9 (64)	35 [2-97]	2 (14)
Warshaw procedure	9 (17)	22 ^e [4-65]	180 [120-348]	555.5 [70-1340]	5 (56)	7 (78)	30 [7-126]	0 (0)
Pancreas-sparing surgery								
Letton–Wilson’s procedure	11 (20)	18 ^c [9-75]	280 [181-690]	514 ^d [0-2575]	3 (27)	7 (64)	31 [7-55]	0 (0)
Bracy procedure	4 (8)	31 [24-83]	398.5 [300-526]	1584.5 [400-2400]	1 (25)	2 (50)	37 [32-52]	0 (0)
No pancreatic resection	5 (10)	32 [19-57]	240 [190-418]	850 [520-11673]	4 (80)	5 (100)	65 ^b [30-175]	0 (0)
Drainage	2 (4)	34.5 [19-50]	304 [190-418]	1,075 [850-1300]	1 (50)	2 (100)	119.5 [64-175]	0 (0)
Pancreatic suture repair	3 (6)	32 [28-57]	240 [221-265]	700 [520-11673]	3 (100)	3 (100)	65 [30-117]	0 (0)

Note: Frequency (%) for categorical variables.

Median (range) for other continuous variables.

^aClavien–Dindo classification of grade IIIa or higher.

^b*P* = .003 vs Pancreatic resection.

^c*P* = .038 vs Pancreaticoduodenectomy.

^d*P* = .010 vs Pancreaticoduodenectomy.

^e*P* = .018 vs Distal pancreatectomy.

mortality increasing with the number of damaged organs.^{3,4} This multicenter nationwide survey conducted over the last decade revealed that mortality was 5% for all cases, 1.8% for patients with isolated pancreatic injury, and 3% for patients with MPD injury. This mortality rate was similar to that reported in previous studies, which found that the mortality rate is 0%-10%.^{2,4,14,15}

Age distribution was analyzed in male and female patients. Young patients were more common regardless of sex. However, male patients showed bimodal distribution with another peak at 60s. Additionally, a similar result was previously reported from Japan.⁴ This might be explained by the fact that males in middle-aged or older populations work outside and drive more frequently compared to the same generation females in this country. In addition, skeletal muscle mass forming visceral protective layer tends to be more vulnerable with aging.

Herman et al. reported that there was no association between elevated serum amylase levels and severity of pancreatic injury according to the American Association for the Surgery of Trauma (AAST) grades, although this indicates that elevated serum amylase levels may be diagnostic

indicators.^{16–18} Alternatively, our analysis was performed between the two groups, focusing on the existence of the MPD injury. Patients with MPD injury had significantly higher serum amylase and CRP levels early after the injury occurred. In this regard, Nadler et al.¹⁹ have also reported that high amylase levels were more consistent with MPD injury. The MPD injury likely contributes to early elevation of serum amylase and CRP. Rapid serum amylase and CRP elevation might be important markers for predicting MPD injury.

Blunt trauma and isolated pancreatic injury were independent factors predicting MPD injury. It is difficult to speculate clear mechanisms for these results. Blunt injury to the pancreas results in the organ being compressed over the vertebra. Localized deep compressive forces may result in an isolated pancreatic laceration depending on the magnitude of the force.²⁰ However, the influence of selection bias in a retrospective setting should be considered.

The diagnostic accuracy of ERP was superior to that of other imaging modalities (CT and MRP) in this survey for the diagnosis of MPD injury. In 1978, Belohravek et al. first reported ERP experience to identify MPD injury in trauma cases.²¹ ERP allows visualization of the MPD and determines

injury severity, facilitating optimal treatment, such as drainage using EPS.⁷ Many other reports also described the value of ERP in patients with pancreatic trauma.^{6-8,11,22} multi-detector-row CT, with sensitivity and specificity of up to 80%, is the best noninvasive imaging modality to detect pancreatic injury.²³ In contrast, although MRP imaging has the advantage of being minimally invasive and can evaluate upstream of the injured area, the rate of implementation is low in Japan because of the lack of examination agility. In patients with suspected deep pancreatic injury on CT examination and stable vital signs, ERP should be performed to evaluate MPD injury more accurately for developing more appropriate treatment strategy.

Complications of the initial treatment were more likely to occur in the MPD injured group, that is, the presence or absence of MPD injury was found to affect short-term outcomes. Regarding which treatment strategy, OM or NOM, should be taken for patients with the MPD injury, OM has been generally recommended.²⁴ Recently published evidence, however, suggested that NOM might be an acceptable and safe option even in patients with MPD injury.⁴ Previous studies have shown that endoscopic management, such as EPS, reduced the need of surgery for pancreatic injury by blocking the pancreatic leak and draining the pancreatic fluid into the duodenum.⁶⁻⁸ Nevertheless, in this survey, NOM was associated with a longer hospital stay than OM. Frequent sequential problems observed in the NOM group most likely resulted in prolonged hospital stay. It is also noteworthy that in this study, five of 11 patients initially treated with NOM subsequently underwent surgery as a secondary treatment although there was no mortality with timely and appropriate additional treatments. In contrast, in a total of 64 patients with MPD injury, six patients eventually received successful NOM and no subsequent surgical intervention. Among them, five patients were treated with endoscopic pancreatic stenting or endoscopic nasopancreatic drainage. NOM including endoscopic stenting procedures may be a less invasive and effective treatment option for selected patients. Because the number of NOM for MPD injury is very small in this series, the actual role of NOM for MPD injury is still unclear and should be explored further in large scale studies.

Our study included a large series of patients with MPD injury who were initially treated with surgery. Pancreatic resection was the more frequently performed procedure in this series. It has been previously reported that emergency PD for pancreatic injury was associated with a very high mortality rate (>30%) because of the patient's poor condition and surgical complexity.^{14,25} However, compared with the outcomes of these previous reports, the outcome of PD in this recent series was excellent with zero mortality. This is presumably

because the attending centers are all JSAEM board-certified hospitals that have abundant experiences in abdominal emergencies. Appropriate initial diagnosis, meticulous perioperative management, prompt decision of surgical intervention based on each patient condition, and recent improvement in perioperative intensive care would be related to the excellent postoperative outcomes. On the other hand, there were only five patients who underwent surgery without pancreatic resection. The length of hospital stay was significantly longer in these five patients than in those who underwent pancreatic resection. Otherwise, pancreatic-conserving surgery including Letton-Wilson and Bracy procedures is preferable to preserve the pancreatic function if the following condition is met: (a) if the patient is young, (b), if the patient's vital signs are stable, and (c) if there is a highly qualified pancreatic surgeon.⁵ In Japan, a surgical qualification system has been implemented by the Japanese Society of Hepatobiliary Pancreatic Surgery.

This study has some limitations as follows. First, the sample size was limited and questionnaires were used to obtain data. Second, because this was a retrospective observational study, selection bias was intrinsic. Third, only board-certified abdominal emergency specialist hospitals of the JSAEM were registered for participation in this survey rather than a large national study. There was an inherent bias in participating institutes. Finally, we were unable to evaluate the long-term outcomes of patients with MPD injury.

In conclusion, serum amylase levels were significantly higher in patients with MPD injury. ERP is the most accurate imaging modality for assessing pancreatic duct injury and is recommended to be performed in patients with stable hemodynamics. Although mortality from pancreatic trauma, including MPD injury, was relatively low regardless of treatment strategy in this series, serious sequelae occurred more frequently and length of hospital stay was significantly longer in patients initially treated nonoperatively.

ACKNOWLEDGMENTS

We are deeply grateful to the Japanese Society for Abdominal Emergency Medicine and the facilities for completing the survey. We also would like to thank Enago (www.enago.com) for English language editing.

CONFLICT OF INTEREST

All authors declare no conflicts of interest.

FUNDING INFORMATION

Project study funds of Japanese Society for Abdominal Emergency Medicine (2016-P-02).

ORCID

Yasuhisa Ando  <https://orcid.org/0000-0002-8686-2810>

Hironobu Suto  <https://orcid.org/0000-0001-8476-3985>

REFERENCES

1. Antonacci N, Di Saverio S, Ciaroni V, Biscardi A, Giugni A, Cancellieri F, et al. Prognosis and treatment of pancreaticoduodenal traumatic injuries: which factors are predictors of outcome? *J Hepatobiliary Pancreat Sci.* 2011;18:195–201.
2. O'Reilly DA, Bouamra O, Kausar A, Malde DJ, Dickson EJ, Lecky F. The epidemiology of and outcome from pancreatoduodenal trauma in the UK, 1989–2013. *Ann R Coll Surg Engl.* 2015;97:125–30.
3. Scollay JM, Yip VS, Garden OJ, Parks RW. A population-based study of pancreatic trauma in Scotland. *World J Surg.* 2006;30:2136–41.
4. Shibahashi K, Sugiyama K, Kuwahara Y, Ishida T, Okura Y, Hamabe Y. Epidemiological state, predictive model for mortality, and optimal management strategy for pancreatic injury: a multicentre nationwide cohort study. *Injury.* 2020;51:59–65.
5. Iacono C, Zicari M, Conci S, Valdegamberi A, De Angelis M, Pedrazzani C, et al. Management of pancreatic trauma: a pancreatic surgeon's point of view. *Pancreatol.* 2016;16:302–8.
6. Kim S, Kim JW, Jung PY, Kwon HY, Shim H, Jang JY, et al. Diagnostic and therapeutic role of endoscopic retrograde pancreatography in the management of traumatic pancreatic duct injury patients: single center experience for 34 years. *Int J Surg.* 2017;42:152–7.
7. Kim HS, Lee DK, Kim IW, Baik SK, Kwon SO, Park JW, et al. The role of endoscopic retrograde pancreatography in the treatment of traumatic pancreatic duct injury. *Gastrointest Endosc.* 2001;54:49–55.
8. Lin BC, Liu NJ, Fang JF, Kao YC. Long-term results of endoscopic stent in the management of blunt major pancreatic duct injury. *Surg Endosc.* 2006;20:1551–5.
9. Dindo D, Demartines N, Clavien PA. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg.* 2004;240:205–13.
10. Bassi C, Dervenis C, Butturini G, Fingerhut A, Yeo C, Izbicki J, et al. Postoperative pancreatic fistula: an international study group (ISGPF) definition. *Surgery.* 2005;138:8–13.
11. Gupta A, Stuhlfaut JW, Fleming KW, Lucey BC, Soto JA. Blunt trauma of the pancreas and biliary tract: a multimodality imaging approach to diagnosis. *Radiographics.* 2004;24:1381–95.
12. Oláh A, Issekutz A, Haulik L, Makay R. Pancreatic transection from blunt abdominal trauma: early versus delayed diagnosis and surgical management. *Dig Surg.* 2003;20:408–14.
13. Potoka DA, Gaines BA, Leppäniemi A, Peitzman AB. Management of blunt pancreatic trauma: what's new? *Eur J Trauma Emerg Surg.* 2015;41:239–50.
14. Kang WS, Park YC, Jo YG, Kim JC. Pancreatic fistula and mortality after surgical management of pancreatic trauma: analysis of 81 consecutive patients during 11 years at a Korean trauma center. *Ann Surg Treat Res.* 2018;95:29–36.
15. Siboni S, Kwon E, Benjamin E, Inaba K, Demetriades D. Isolated blunt pancreatic trauma: a benign injury? *J Trauma Acute Care Surg.* 2016;81:855–9.
16. Moore EE, Cogbill TH, Malangoni MA, Jurkovich GJ, Champion HR, Gennarelli TA, et al. Organ injury scaling, II: pancreas, duodenum, small bowel, colon, and rectum. *J Trauma.* 1990;30:1427–9.
17. Herman R, Guire KE, Burd RS, Mooney DP, Ehlich PF. Utility of amylase and lipase as predictors of grade of injury or outcomes in pediatric patients with pancreatic trauma. *J Pediatr Surg.* 2011;46:923–6.
18. Mahajan A, Kadavigere R, Sripathi S, Rodrigues GS, Rao VR, Koteswar P. Utility of serum pancreatic enzyme levels in diagnosing blunt trauma to the pancreas: a prospective study with systematic review. *Injury.* 2014;45:1384–93.
19. Nadler EP, Gardner M, Schall LC, Lynch JM, Ford HR. Management of blunt pancreatic injury in children. *J Trauma.* 1999;47:1098–103.
20. Antonsen I, Berle V, Søreide K. Blunt pancreatic injury in children. *Tidsskr Nor Lægeforen.* 2017;137:17–28.
21. Belohlavek D, Merkle P, Probst M. Identification of traumatic rupture of the pancreatic duct by endoscopic retrograde pancreatography. *Gastrointest Endosc.* 1978;24:255–6.
22. Kumar A, Panda A, Gamanagatti S. Blunt pancreatic trauma: a persistent diagnostic conundrum? *World J Radiol.* 2016;8:159–73.
23. Degiannis E, Glapa M, Loukogeorgakis SP, Smith MD. Management of pancreatic trauma. *Injury.* 2008;39:21–9.
24. Lin BC, Wong YC, Chen RJ, Liu NJ, Wu CH, Hwang TL, et al. Major pancreatic duct continuity is the crucial determinant in the management of blunt pancreatic injury: a pancreatographic classification. *Surg Endosc.* 2017;31:4201–10.
25. van der Wilden GM, Yeh D, Hwabejire JO, Klein EN, Fagenholz PJ, King DR, et al. Trauma Whipple: do or don't after severe pancreaticoduodenal injuries? An analysis of the National Trauma Data Bank (NTDB). *World J Surg.* 2014;38:335–40.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section.

How to cite this article: Ando Y, Okano K, Yasumatsu H, et al. Current status and management of pancreatic trauma with main pancreatic duct injury: A multicenter nationwide survey in Japan. *J Hepatobiliary Pancreat Sci.* 2021;00:1–9. <https://doi.org/10.1002/jhbp.877>

